



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Region and task-specific activation of Arc in primary motor cortex of rats following motor skill learning

Hosp, J A ; Mann, S ; Wegenast-Braun, B M ; Calhoun, M E ; Luft, A R

Abstract: Motor learning requires protein synthesis within the primary motor cortex (M1). Here, we show that the immediate early gene Arc/Arg3.1 is specifically induced in M1 by learning a motor skill. Arc mRNA was quantified using a fluorescent in situ hybridization assay in adult Long-Evans rats learning a skilled reaching task (SRT), in rats performing reaching-like forelimb movement without learning (ACT) and in rats that were trained in the operant but not the motor elements of the task (controls). Apart from M1, Arc expression was assessed within the rostral motor area (RMA), primary somatosensory cortex (S1), striatum (ST) and cerebellum. In SRT animals, Arc mRNA levels in M1 contralateral to the trained limb were 31% higher than ipsilateral ($p < 0.001$), 31% higher than in the contralateral M1 of ACT animals ($p < 0.001$) and 48% higher than in controls ($p < 0.001$). Arc mRNA expression in SRT was positively correlated with learning success between two sessions ($r = 0.52$; $p = 0.026$). For RMA, S1, ST or cerebellum no significant differences in Arc mRNA expression were found between hemispheres or across behaviors. As Arc expression has been related to different forms of cellular plasticity, these findings suggest a link between M1 Arc expression and motor skill learning in rats.

DOI: <https://doi.org/10.1016/j.neuroscience.2013.06.060>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-84738>

Journal Article

Accepted Version

Originally published at:

Hosp, J A; Mann, S; Wegenast-Braun, B M; Calhoun, M E; Luft, A R (2013). Region and task-specific activation of Arc in primary motor cortex of rats following motor skill learning. *Neuroscience*, 250:557-564.

DOI: <https://doi.org/10.1016/j.neuroscience.2013.06.060>

Region and task-specific activation of Arc in primary motor cortex of rats following motor skill learning

Jonas A. Hosp ^{1,2}, Sarah Mann ^{1,2}, Bettina M. Wegenast-Braun ³, Michael E. Calhoun ³, Andreas R. Luft ^{1,2,4,5}

Formatiert: Deutsch (Schweiz)

¹ Clinical Neurorehabilitation, Department of Neurology, University of Zurich, Zurich, Switzerland

Formatiert: Englisch (USA)

² Rehabilitation Initiative and Technology Center (RITZ), Zurich

³ Department of Cellular Neurology, Hertie Institute for Clinical Brain Research, and DZNE, German Center for Neurodegenerative Diseases, Tübingen, Germany

⁴ Division of Brain Injury Outcomes, Department of Neurology, Johns Hopkins University, Baltimore, MD, USA

⁵ Center for Integrative Human Physiology, University of Zurich, Switzerland

Formatiert: Englisch (USA)

Running Head: Arc induction by motor learning

Correspondence to:

Andreas R. Luft
Department of Neurology
University of Zurich
Frauenklinikstrasse 26
8091 Zurich
Switzerland
Tel +41 44 255 5400
Fax +41 44 255 4649

Formatiert: Deutsch (Schweiz)

andreas.luft@usz.ch

Formatiert: Schriftart: Cambria,
Nicht unterstrichen, Schriftartfarbe:
Schwarz

Formatiert: Englisch (USA)

Abstract

~~Motor~~ Motor learning requires protein synthesis within primary motor cortex (M1).

Formatiert: Englisch (USA)

Here, we show that the immediate early gene *Arc/Arg3.1* is specifically induced in M1 by learning a motor skill.

Arc mRNA was quantified using a fluorescent in situ hybridization assay in adult Long-Evans rats learning a skilled reaching task (SRT), in rats performing reaching-like forelimb movement without learning (ACT) and in rats that were trained in the operant but not the motor elements of the task (controls). Apart from M1, *Arc* expression was assessed within the rostral motor area (RMA), primary somatosensory cortex (S1), striatum (ST) and cerebellum.

In SRT animals, *Arc* mRNA levels in M1 contralateral to the trained limb were 31% higher than ipsilaterally (p<0.001), 31% higher than in the contralateral M1 of ACT animals (p<0.001) and 48% higher than in controls (p<0.001). *Arc* mRNA expression in SRT was positively correlated with learning success between two

sessions ($r=0.52$; $p=0.026$). For RMA, S1, ST or cerebellum no significant differences in *Arc* mRNA expression were found between hemispheres or across behaviors.

As *Arc* expression has been related to different forms of cellular plasticity, these findings suggest a link between M1 *Arc* expression and motor skill learning in rats.

~~learning requires protein synthesis within primary motor cortex (M1). The proteins expressed to mediate successful learning are largely unknown. Here, we show that the immediate early gene *Arc/Arg3.1* is specifically induced within the caudal motor cortex (M1) by learning a motor skill.~~

~~*Arc* mRNA was quantified using a fluorescent in situ hybridization assay (FISH) in adult Long-Evans rats learning a skilled reaching task (SRT), in rats performing reaching-like forelimb movement without learning (activity control task, ACT) and in rats that were trained in the operant but not the motor elements of the task (control group). Apart from M1, *Arc* expression was also assessed within the rostral motor area (RMA), primary somatosensory cortex (S1), striatum (ST) and cerebellum.~~

~~In SRT animals, *Arc* mRNA levels in M1 contralateral to the trained limb were 31% higher ipsilaterally ($p<0.001$), 31% higher than in the contralateral M1 of ACT animals ($p<0.001$) and 48% higher than in control animals ($p<0.001$). *Arc* mRNA expression in SRT was positively correlated with learning success between two sessions ($r=0.52$; $p=0.026$). For RMA, S1 and ST, no significant~~

differences in *Arc* mRNA expression were found between hemispheres or across behaviors. For cerebellum, only negligible low levels of *Arc* mRNA were observed. As *Arc* expression has been related to different forms of cellular plasticity, these findings suggest a link between M1 *Arc* expression and motor skill learning in rats.

Key words: Arc, FISH, motor learning, motor cortex

Formatiert: Englisch (USA)

Formatiert: Block, Abstand Nach:
10 Pt., Zeilenabstand: Doppelt

Formatiert: Abstand Nach: 10 Pt.

Formatiert: Englisch (USA)

Introduction

While the process of motor skill learning depends on the interaction of different brain regions (e.g. ~~namely~~ sensorimotor cortex, (SM), basal ganglia, (BG) and cerebellum, (C; Hikosaka et al., 2002), evidence points to primary motor cortex (M1) as the structure where motor memory traces are formed. Skill acquisition requires protein synthesis within M1 and induces long lasting changes in M1 in synaptic strength (Riout-Pedotti et al., 2000), ~~in M1 possibly reflecting the~~ storage mechanisms for motor memories (Kleim et al., 2003, Luft et al., 2004). ~~However, Learning induces long lasting changes in M1 synaptic strength (Riout-Pedotti et al., 2000).~~ Little is known about the genes and proteins that mediate these processes.

In rats, the immediate-early gene (IEG) c-fos is expressed within M1 after beginning to training an acrobatic locomotor skill, ~~and C-fos~~ remains elevated when a performance plateau has been reached (Kleim et al., 1996). Recently, increased levels of Arc (activity-regulated cytoskeleton-associated protein), a protein coded by the IEG Arc (also known as the activity-regulated gene 3.1 Arg3.1) have been found in M1 of rats that trained precision reaching task with the contralateral forelimb (Hanlon et al., 2009). As this study focused on the

Formatiert: Block, Abstand Nach: 10 Pt., Zeilenabstand: Doppelt

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert ...

Formatiert: Hervorheben

Formatiert ...

Formatiert: Hervorheben

Formatiert ...

Formatiert: Englisch (USA)

Formatiert ...

Formatiert: Hervorheben

Formatiert ...

Formatiert ...

Formatiert: Hervorheben

Formatiert ...

Formatiert: Englisch (USA)

Formatiert ...

effect of motor training on ~~slow-wave activity an cortical~~ IEG expression during non-REM sleep, ~~its an open question if it did not reveal whether~~ Arc-induction was specific to learning or simply related to ~~e~~-activity, i.e. moving the forelimb more than usually.

Formatiert: Hervorheben

Formatiert: Englisch (USA),
Hervorheben

Formatiert: Englisch (USA),
Hervorheben

As IEGs have been extensively studied within the hippocampal network, all knowledge summarized here was obtained from this system unless cited differently. IEGs typically are transcribed within few minutes after induction

Formatiert: Hervorheben

of long-term potentiation (LTP, Worley et al., 1993; Guzowski et al., 1999). In contrast to the IEG *c-fos*, an activity-induced transcription factor that controls the

Formatiert: Englisch (USA),
Hervorheben

Formatiert: Hervorheben

expression of other transcription factors, *Arc* is an "effector-IEG" ~~that that~~

Formatiert: Hervorheben

~~promotes induces the~~ transcription of proteins ~~directly related to cellular modifications influencing such as changes in the~~ cytoskeleton or synaptic AMPA

Formatiert: Hervorheben

Formatiert: Hervorheben

receptor trafficking (Bramham et al., 2008, Miyashita et al., 2008). These modifications are thought to ~~mediate learning related underly~~ cellular plasticity.

Formatiert: Englisch (USA),
Hervorheben

Formatiert: Hervorheben

Learning and experience-related transcription of *Arc* mRNA has been observed in various behavioral paradigms such as the Morris water maze task (Guzowski

Formatiert: Englisch (USA),
Hervorheben

et al., 2000, Fletcher et al., 2006). In animals not subjected to learning

paradigms, *Arc* is transcribed at very low levels. Upon excitatory synaptic

activation *Arc* is expressed within minutes (Lyford et al., 1995) in an "all-or-

Formatiert: Englisch (USA)

nothing" fashion (Guzowski et al., 1999). Its induction is confined to neural

assemblies associated with the encoding of information of specific behavioral

experiences (Steward et al., 1998, Guzowski et al., 1999). After induction, *Arc*

mRNA is transported into dendrites and accumulates at sites of synaptic

activation where it is locally translated into proteins (Steward et al., 1998). Hence, Arc can be considered as a cellular marker of learning-related synaptic plasticity.

The objective here was to investigate, if whether M1 Arc induction is specific for motor skill learning or whether it is related to motor activity in the absence of learning. Further, we investigated whether Arc becomes, is induced in M1 and other brain regions related to motor learning in a concerned with learning-specific manner, motor skill learning and if whether the degree of Arc induction is related to learning efficacy.

Formatiert: Englisch (USA)

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA)

Material and Methods

Animals

Adult male Long-Evan rats (8-10 weeks, 250-350 g, Centre d'Elevage R. Janvier, Le Genest-St. Isle, France) were used for all experiments. Animals were housed individually in a 12/12-hr light/dark cycle (light on: 8pm, off: 8am). Littermates were distributed equally among the groups of an experiment. All experiments were conducted in accordance with German and Swiss regulations and were approved either by the Animal Commission of the State of Baden-Württemberg or the Committee for Animal Experimentation of the Canton of Zürich.

Behavioral conditions

Training sessions were performed at the beginning of the dark phase. Animals were food-restricted for 24 hr before the first training session. During training animals were kept slightly over their initial weight by providing 50 mg/kg of standard lab diet after each training session. Water was given *ad libitum*. The behavioral tasks were performed as previously described (Molina-Luna et al., 2008). The training cage was a 15 x 40 cm chamber (height 30 cm) with a vertical window (1 cm wide, 5 cm high, lower edge 2 cm above ground) in the front wall and a small light sensor in the rear wall (7 cm above ground).

Three different behavioral conditions were compared: a motor skill learning paradigm (skilled reaching task; SRT), ~~and~~ a paradigm requesting arm movements without motor learning (activity control task; ACT) and controls with the operant but without the motor elements (control group; CG). ~~Animals were killed after pre-training were used as a control.~~ These rats were exposed to the same training cage and had accessed a food pellet by tongue (pre-training).

Because for technical reasons animals from the three groups could be evaluated in the same immunohistochemistry run, pairwise matching was performed between SRT/ACT and SRT/~~control~~ in two runs. Animals in the SRT/ACT or SRT/CG pairings were trained for exactly the same amount of time.

As behavioural tasks were embedded in an operant conditioning paradigm, animals required a pre-training to operate the experimental setup properly, before being assigned to a particular experimental group. During this pre-training, animals learned to open the motorized sliding door that covered the front window by nose-poking the sensor in the rear. Opening the window gave access

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftartfarbe: Automatisch, Englisch (USA), Hervorheben

Formatiert: Schriftart: 12 Pt., Schriftartfarbe: Automatisch, Englisch (USA), Hervorheben

Formatiert: Schriftartfarbe: Automatisch, Englisch (USA), Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Englisch (USA), Hervorheben

to one food pellet (45 mg, Bio-serve, Frenchtown, NJ, USA) located on a small horizontal board in a distance of 0.5 cm relative to the outside edge of the window. Thus, pellets could be retrieved. During pre-training, pellets were retrieved by tongue without utilization of the forepaw. Upon retrieval a pellet dispenser automatically replaced the pellet. Whereas SRT and ACT animals were pre-trained for five days, control animals were killed after two days of pre-training.

Skilled reaching task (SRT):

In SRT animals pre-training was followed by motor training that was initiated by removing the board and placing the pellet on a small vertical post 1.5 cm away from the window. In this position pellets were only retrievable by using the forelimb. Because the diameter of the post was approximately that of the pellet, the pellet was in an unstable position easily kicked off the post. Before the first skill training session, forelimb preference was determined. Then the pedestal was shifted to one side of the window to allow for reaching with the preferred limb only. To retrieve the pellet rats had to extend the forelimb towards the target, pronate, open the paw, grasp, and pull the forelimb back while supinating to bring the pellet towards the mouth (Whishaw and Pellis, 1990). Each reaching trial was scored as "successful" (reach, grasp and retrieve) or "unsuccessful" (pellet pushed off pedestal or dropped during retraction).

For the skilled-reaching task in male Long-Evans rats, motor learning seems to be especially effective during the second day of training as the highest increase in learning success occurs between sessions two and three (Buitrago et al.).

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA), Hervorheben

Formatiert: Englisch (USA)

Formatiert: Schriftartfarbe: Rot, Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftartfarbe: Automatisch, Englisch (USA), Hervorheben

2004). Regarding the intra-session learning curve at day two, the steepest increase in successful grasps can be found between trials 40 to 60 (Buitrago et al., 2004). To display expression of *Arc* mRNA at this particular sensitive time-point, animals were killed after The first session (d1) consisted of 100 door openings (= trials), the second session (d2) consisted of only 50 trials at day 2 whereas training day 1 consisted of 100 trials. to focus on the time point when the averaged learning curve has its steepest course (Buitrago et al., 2004). The

Formatiert: Hervorheben

Formatiert: Englisch (USA)

improvement of reaching performance between sessions was defined as the difference of successful trials between training on day 2 (50 trials) and – to render the comparison valid – the first half (50 trials) of the training session on day 1. In case rats showed a lower performance at day 2 compared to day 1, negative values of learning rates were depicted.

Activity control task (ACT)

The ACT consisted of extending the forelimb through the window to touch a sensor in 1.5 cm distance. If the sensor was touched, the investigator gave the rat a pellet directly into the mouth of the rat using forceps. Limb position during reaching in ACT was identical to SRT but no grasping or pellet retrieval was necessary.

Formatiert: Englisch (USA)

Tissue preparation

Are mRNA can be detected in the cytoplasm of neurons about 20 to 45 minutes after induction (Guzowski, 2002). The training session on day 2 lasted approximately 20 minutes. Animals were killed 15 minutes after the second training session ended. Animals were then deeply anesthetized (pentobarbital; 50 mg/kg i.p.; Kantonsapotheke Zurich, Switzerland) and perfused transcardially with PBS followed by 4% paraformaldehyde (PFA). All solutions used for animal dissection were prepared RNase-free. The brains were removed and kept for 24 hours in 4% PFA, then for 24 hours in 10% glycerol and finally for 1-2 days in 20% glycerol all at 4°C. Subsequently, brains were rapidly frozen in 2-methylbutane. Coronal sections (40µm) were prepared using a freezing/sliding microtome (Leica Microsystems GmbH, Wetzlar, Germany). For each animal, three brain sections containing the forelimb representation of M1 (3 mm ant. to 0.5 mm anterior to bregma), RMA (3.5 mm ant. to 1.5 mm anterior to bregma), S1 (1 mm ant. to 1.5 mm posterior to bregma) and striatum (2 mm anterior to 0.5 mm posterior to bregma) were collected, stereotactic coordinates were derived from Paxinos and Watson, 1998. The three sections were equally spaced (fixed distance of 600µm between subsequent slices) with a randomly chosen starting point, which was retained for matched behavioral pairings. In principle, a systematic random sampling approach (random starting point, fixed distance of 600µm between subsequent slices) was applied for selection of brain slices. However, to remain comparability between groups, similar starting points were used for matched pairs. For SRT and ACT animals, both hemispheres were sampled. The hemisphere contralateral to the forelimb preferred for reaching will

Formatiert: Hervorheben

Formatiert: Kommentartext, Absatzkontrolle, Tabstopps: Nicht an 0.99 cm + 1.98 cm + 2.96 cm + 3.95 cm + 4.94 cm + 5.93 cm + 6.91 cm + 7.9 cm + 8.89 cm + 9.88 cm + 10.86 cm + 11.85 cm

Kommentar [M1]: Waren das 3 Sections je Region, oder 3 Sections die alle Regionen beinhaltet haben? – ist momentan nicht klar.

Formatiert: Nicht Hervorheben

Formatiert: Schriftart: (Standard) Helvetica

Formatiert: Nicht Hervorheben

Formatiert: Schriftart: (Standard) Helvetica, Hervorheben

Formatiert: Nicht Hervorheben

Kommentar [M2]: Strenggenommen hätte man dann eigentlich jeden 15. Schnitt verwenden sollen und nicht auf 3 Schnitte begrenzen.

Formatiert: Nicht Hervorheben

Kommentar [M3]: Würde man stereologisch auch nicht so machen.

Mein Vorschlag unten versucht die stereologischen Schwachpunkte etwas zu kaschieren.

Kommentar [M4]: The three sections were equally spaced (fixed distance of 600µm between subsequent slices) with a randomly chosen starting point, which was retained for matched behavioral pairings.

Formatiert: Nicht Hervorheben

be referred to as “contralateral hemisphere” (cSRT and cACT), the opposite hemisphere as ipsilateral (iSRT and iACT).

Formatiert: Englisch (USA)

Arc probe generation

Riboprobes were generated from nearly full-length rat *Arc* cDNA inserted into pBluescript II SK plasmid (provided by Drs. J. Guzowski and P. Worley). Sense and antisense digoxigenin-labeled probes were transcribed from the linearized plasmid using DIG RNA Labeling Mix, and T3 or T7 RNA polymerases respectively (Stratagene, La Jolla, CA). Probes were then purified on mini Quick Spin RNA Columns. ~~To improve tissue penetration, probes were alkaline hydrolyzed for 30 minutes at 60°C.~~ Probe concentration was estimated by spotting serial dilutions of probe and DIG-labeled control RNA on membranes, immunodetection with anti-DIG Fab fragments conjugated to alkaline phosphatase, and colorization with nitro blue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate. All reagents were obtained from Roche Diagnostics, Mannheim, Germany unless indicated otherwise.

Formatiert: Englisch (USA)

In situ hybridization immunohistochemistry

Fluorescent *in situ* hybridization was performed using a modified version of the protocol described by Wegenast-Braun et al., 2009, allowing for application in rats. Solutions used for *in situ* hybridization were prepared RNase-free. Sections were mounted onto slides (SuperFrostPlus; R. Langenbrinck Labor- und Medizintechnik, Teningen, Germany). If mounted sections were processed

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

individually, differences in staining intensities could be a possible source of confounds. To overcome this problem, brain sections from different experimental groups were distributed in a balanced fashion, i.e. sections from each brain region derived from a SRT animal were mounted pairwise with their counterparts from the ACT or control group on the same slide. Aside from this rule, pairs of sections were distributed in a random order.

Mounted slides were dried for 1 hour at 30°C, and postfixed for 20 minutes in 4% paraformaldehyde in PBS at 4°C, followed by three washes in PBS. Sections were permeabilized with 0.3% Triton-X100 in PBS and 30 µg/ml proteinase K in Tris EDTA (TE) [10 mM Tris-HCl and 1 mM EDTA, pH 7.5] at 30°C with interleaved PBS washes. Proteinase K treatment was stopped by 0.2% glycine in PBS with 50 mM EDTA followed by two PBS washes. Tissue was then equilibrated in 0.1 M triethanolamine (TEA) and acetylated with 0.25% acetic anhydride in 0.1 M TEA. After three washes in 2x SSC [0.3 M NaCl and 0.03 M sodium citrate, pH adjusted to 7.0 with citric acid], sections were dehydrated through graded alcohols and defatted in chloroform. Tissue was rehydrated through graded alcohols and dried. For hybridization, the probe RNA was diluted to 100 ng/ml in hybridization buffer [50% deionized formamide, 2x SSC, 1 Denhardt's solution (1% bovine serum albumin, 1% polyvinylpyrrolidone, and 1% Ficoll), 10% dextran sulfate, 0.5 mg/ml wheat germ tRNA, and 0.5 mg/ml salmon sperm DNA] and denatured at 80°C for 5 minutes before application to tissue. Sections were then sealed with a coverslip and incubated 16 to 20 hours at 59°C in a humid chamber. Coverslips were removed and slides were washed twice in

2x SSC/1 mM EDTA. Then, sections were treated with 10 µg/ml RNase A (Sigma-Aldrich, Steinheim, Germany) in TE/0.5 M NaCl and washed twice with 2x SSC/1 mM EDTA. Stringency washes were performed in 0.1x SSC/1 mM EDTA at 59°C for 1-2 hours and two times in 0.5 SSC at room temperature for 5 minutes. Endogenous peroxidase activity was quenched with 1% H₂O₂ in PBS, followed by subsequent washes in PBS, washing buffer [100 mM maleic acid, 150 mM sodium chloride, 0.3% Tween-20, pH adjusted to 7.5] and tris-buffered saline (TBS) (84 mM Tris-HCl, 16 mM Trisbase, and 154 mM NaCl). Sections were blocked with 10% bovine serum albumin in TBS and incubated with anti-DIG Fab fragments conjugated to horseradish peroxidase diluted 1:400 in 10% bovine serum albumin in TBS for 2 hours at 4°C for detection. After three washes with washing buffer, the conjugate was detected with Cyanine-3 Tyramide Working Solution (TSAPlus Fluorescence Systems, Perkin Elmer). Developed sections were washed with TE and PBS, counterstained with 4'-6-diamidino-2-phenylindole (DAPI; Sigma-Aldrich) and allowed to dry before adding coverslips with VectaShield (Vector Laboratories, Burlingame, CA

✂

Confocal microscopy and data analysis

Regions of interests (ROIs) were identified by cortical cytoarchitecture: S1 could be easily identified due to prominent layer IV, M1 on the other hand by the absence of such a granular layer. RMA could be defined due to high neuronal

Formatiert: Englisch (USA)

Formatiert: Hervorheben

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Hervorheben

Formatiert: Abstand Nach: 0 Pt.

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Hervorheben

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Hervorheben

density and a feathering of the strict layered order of cortex towards midline.

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Hervorheben

Mosaic-images of the particular ROIs comprising the ROI-entire anatomic region

Formatiert: Hervorheben

of interest (e.g. M1) were taken with a Zeiss LSM 510 META (Axiovert 200M)

confocal microscope equipped with three lasers (488, 543 and 633 nm) using a

Zeiss Plan-Neofluar 20x/0.5 air objective. Settings were kept constant for

Formatiert: Nicht Hervorheben

different imaging sessions. Laser powers were regularly measured with a PT

9610 laser power analyzer (Gigahertz-Optik, Puchheim, Germany) to control for

changes in intensities. To avoid edge artifacts at the cut surfaces of the tissue,

the optical focal plane was set below the surface, well within the penetration

margin of the in situ hybridization signal. with the size The optical thickness of

was set to 5.6 μm was taken from the middle of the brain slices.

Formatiert: Nicht Hervorheben

Formatiert: Nicht Hervorheben

To avoid an oversampling by including Arc mRNA positive neurons of

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Hervorheben

neighboring structures, the counting area (ROI) was set with a lateral security

Kommentar [M5]: manually ?

zone of 50 μm apart from the microscopically defined border. RGB images were

Formatiert: Hervorheben

analyzed offline using ImageJ software, (Cell counter plugin) by hand counting

Formatiert: Hervorheben

neurons within this marked area showing expression of Arc mRNA (Cy3) within

Formatiert: Hervorheben

one region of interest. Training sessions on day 2 had a duration of

Formatiert: Hervorheben

Formatiert: Hervorheben

Kommentar [M6]: for each

Kommentar [M7]: Übergang?

approximately 20 minutes and animals were killed exactly 15 minutes after the

Formatiert: Hervorheben

end of the behavioural training. Arc mRNA is present in the cytoplasm of neurons

Formatiert: Nicht Hervorheben

20 to 45 minutes after its induction (Guzowski, 2002), only cells with a clear

Formatiert: Nicht Hervorheben

cytoplasmic staining-pattern were counted to ensure that only the Arc mRNA

Formatiert: Schriftart: Kursiv

Formatiert: Hervorheben

expression induced by the recent training session was measured.

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Englisch (USA), Hervorheben

Although the sampling and counting strategy used here is in good accordance

Formatiert: Standard, Abstand
Nach: 0 Pt., Tabstops: Nicht an
0.99 cm + 1.98 cm + 2.96 cm +
3.95 cm + 4.94 cm + 5.93 cm +
6.91 cm + 7.9 cm + 8.89 cm +
9.88 cm + 10.86 cm + 11.85 cm

with stereological rules, due to the lower number of sections sampled for each subregion (3), we resigned to extrapolate a total number of stained neurons per region and decided to present an average number of activated neurons per counted section within a region. Average values for number of activated neurons were summed across the three analyzed sections for each brain region. Because the overall number of samples was large, two runs of *Arc* mRNA in situ hybridization were required (SRT animals n=6 + controls n=6; SRT animals n=12 + ACT animals n=12). To account for potential differences in effectiveness and intensity of *Arc* mRNA-staining between hybridization runs, the numbers of positive cells of ACT and control group were normalized to the average value of their respective SRT group. Statistica version 7.0 (StatSoft, Inc., Tulsa, OK, USA) software was used to perform ANOVA tests for each brain region. Group and hemisphere were entered as independent variables. The dependent variable was the normalized number of *Arc* mRNA-positive neurons. If the overall ANOVA was significant, Bonferroni post hoc tests were computed. The relationship between *Arc* expression and inter-session improvement was assessed by linear regression analysis. Numerical results are expressed as mean and \pm SEM.

Formatiert: Englisch (USA)

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Englisch (USA), Hervorheben

Formatiert: Englisch (USA)

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Englisch (USA), Hervorheben

Formatiert: Schriftart: 12 Pt.,
Schriftartfarbe: Automatisch,
Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA), Nicht
Hervorheben

Kommentar [M8]: Sind die SRT
Tiere unabhängig, insgesamt also 18?

Formatiert: Englisch (USA)

Formatiert: Englisch (USA), Nicht
Hervorheben

Kommentar [M9]: Also different
behavioral runs?

Formatiert: Englisch (USA)

Formatiert: Englisch (USA), Nicht
Hervorheben

Kommentar [M10]: Waren die
Werte sehr unterschiedlich?
Andernfalls könnte man sie sogar
benennen?

Formatiert: Englisch (USA)

Formatiert: Englisch (USA), Nicht
Hervorheben

Formatiert: Englisch (USA)

Results

In M1 *Arc* expression was differentially affected by the type of training and the hemisphere (ipsilateral versus contralateral; $F_{4,60}=33.7$ $p<0.001$). Post-hoc analysis revealed that the contralateral (trained) hemisphere of SRT (cSRT) animals showed stronger of *Arc* mRNA expression when compared to the ipsilateral hemisphere (iSRT, had 69% of *Arc* mRNA positive neurons compared to cSRT; $p<0.001$), cACT (had 69% of *Arc* mRNA positive neurons compared to cSRT; $p<0.001$), iACT (had 66% of *Arc* mRNA positive neurons compared to cSRT; $p<0.001$) and control group (had 52% of *Arc* mRNA positive neurons compared to cSRT; $p<0.001$; **Fig. 1a**). The number of *Arc* mRNA positive neurons within M1 of the control animals was significantly smaller when compared to the iSRT ($p=0.019$) and cACT group ($p=0.037$; **Fig. 1a**). *Arc* mRNA positive neurons were observed within all cortical layers within M1 except layer I (**Fig. 1b and c**).

The number of *Arc* mRNA positive neurons within M1 of the cSRT group was positively correlated to the learning rate (R-value: 0.523, $p=0.026$; **Fig. 2**), i.e. magnitude of improvement in grasping performance (d1 vs. d2) is related to a

higher Arc expression in M1 contralateral to the trained paw. Furthermore, there is also a non-significant trend towards a correlation between *Arc* mRNA expression in M1 of the ipsilateral hemisphere (iSRT) and learning rate (R-value: 0.402, $p=0.098$).

For RMA ($F_{4,56}=1.0$, $p=0.394$; **Fig. 3a**), S1 ($F_{4,49}=0.5$, $p=0.7$; **Fig. 3b**) and ST ($F_{4,54}=1.258$, $p=0.298$; **Fig 3c**), no significant differences in *Arc* mRNA expression could be detected across behaviors. The number of Arc RNA-positive neurons in cerebellum was very low in all conditions, and was not quantitatively analyzed.

Discussion

This study demonstrates that *Arc* is induced in M1 shortly after motor skill acquisition at a significantly higher level than following motor performance without related learning. The degree of *Arc* expression correlates with performance, i.e. how successfully the motor skill was acquired.

Activation of *Arc* in neurons has been studied in various learning and conditioning paradigms. For example, *Arc* is induced in the olfactory cortex after exposure to odorants (Zou et al., 2005) or in hippocampus after spatial exploratory behavior (Ramirez-Amaya et al., 2005, Fletcher et al., 2006). Moreover, *Arc* is not only a marker of cellular activity related to learning but also seems to be essential for memory consolidation in hippocampus-dependent

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

learning. Infusing *Arc* antisense desoxyribonucleic acids (ODNs) into the hippocampus impaired the formation of spatial memory in the water maze (Guzowski et al., 2000) and the retention performance in an inhibitory avoidance task (McIntyre et al., 2005). *Arc* knockout mice fail to generate long-lasting memory in implicit and explicit learning tasks (Plath et al., 2006).

Here, the expression of *Arc* induced by different learning paradigms was studied in brain regions associated with motor control. *Arc* expressing neurons could be frequently detected in all brain regions that were sampled (M1, RMA, S1 and ST) except in cerebellum. This confirms previous studies showing *Arc* induction in α -calcium-calmodulin kinase II (α CaMKII) positive neurons in hippocampus, neocortex or striatum (Vazdarjanova et al., 2006), whereas cerebellar Purkinje cells are known to show minimal expression (Smith-Hicks et al., 2010). RMA, S1 and ST showed no statistically significant differences in *Arc* expression between different learning conditions or hemispheres. *Arc* was significantly upregulated selectively within M1 contralateral to the trained paw (cSRT) as compared to the ipsilateral hemisphere (iSRT) or to M1 of ACT and control animals. The degree of *Arc* induction correlated with learning success, suggesting a mechanistic link (without proving causation). These findings give further support to the assumption that M1 is an area where motor memories are consolidated (Luft et al., 2004).

Previous studies showing that *Arc* inhibition by antisense desoxyribonucleic acids (ODNs) impaired memory formation in hippocampus dependent learning paradigms (Guzowski et al., 2000, McIntyre et al., 2005, Plath et al., 2006) suggest that similar inactivation in M1 would also impair motor

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Feldfunktion geändert

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen, Hervorheben

Formatiert: Hervorheben

~~learning but further studies are necessary to prove this hypothesis.~~

Hanlon et al. (2009) reported Arc expression in layer V neurons of M1 after rats were trained in reaching (Hanlon et al., 2009). This study was focused on the effect of motor training on slow wave activity and cortical IEG expression during non-REM sleep. ~~Because immunohistochemistry against Arc protein was used,~~ the findings had to be interpreted with caution. ~~Immunohistochemistry measures Arc protein induction that does not follow as predictable a time course following learning. In contrast to protein induction, in contrast, transcription production and~~ intracellular transport of Arc mRNA, ~~which was measured by FISH here, has~~ a well-defined time course (Guzowski et al., 1999, Guzowski, 2002) and, hence, can be unequivocally linked to the learning that occurred shortly before. By measuring cytoplasmic mRNA, we therefore ensured that only those neurons in which Arc was induced by the motor training were visualized. Also, Hanlon and colleagues (2009) did not study control groups and, hence, cannot demonstrate learning-specific Arc induction.

In contrast to the findings of Hanlon et al. we found Arc mRNA to be expressed across all cortical layers except layer I. ~~Arc mRNA is translated locally at active synaptic sites (Steward et al., 1998). Its localization and accumulation there can be enhanced by activating the extracellular signal-regulated kinase (ERK; Huang et al., 2007). Thus, as Furthermore, in rat dentate gyrus granule cells, ERK activation enhances the phosphorylation of the eucaryotic initiation factor 4E (eIF4E) resulting in an increased formation of translationally competent ribosomes and increased synthesis of Arc protein. (Richter and Sonenberg,~~

Formatiert: Englisch (USA)

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Englisch (USA),
Hervorheben

Formatiert: Hervorheben

Formatiert: Schriftart: Cambria,
Schriftartfarbe: Schwarz,
Rechtschreibung und Grammatik
prüfen, Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

Formatiert: Hervorheben

2005, Panja et al., 2009). Thus, Arc mRNA trafficking and as well as its local translation are highly regulated processes that depend on local cofactors like activation of the ERK pathway. Therefore, it cannot be assumed that nuclear Arc mRNA expression and local synaptic Arc protein content are directly proportional.

Formatiert: Hervorheben

Induction of Arc mRNA that occurs within minutes after a neuron became activated in relation to behavioral experience (Guzowski et al., 1999) seems to be an early step specifically tied to the encoding of a motor memory in M1. However, subsequent cascades of gene expression and maturation processes likely have to take place until the memory is fully consolidated (Luft and Buitrago, 2005). Arc mRNA trafficking, local translation of Arc mRNA into protein as well as the turnover of Arc mRNA is a strictly regulated process (Giorgi et al., 2007, Huang et al., 2007, Park et al., 2008, Bramham et al., 2010). Thus, the broad expression pattern of Arc mRNA comprising all cortical layers of M1 (except layer I) (Fig. 1b and c) may reflect an initial activation of neuronal populations in response to training. Later different processes may be required to selectively stabilize newly formed synapses (Xu et al., 2009) or strengthen specific synaptic connections that are functionally relevant.

Similar to Arc here, enhanced expression of the IEG *c-fos* in rat M1 occurred during acrobatic skill training (Kleim et al., 1996). As *c-fos* expression was highest during skill acquisition and decreased during maintenance phase, the learning-specificity of IEG expression in M1 is highly plausible. But besides the upregulation of Arc expression within M1 of the trained hemisphere in SRT rats, numbers of Arc mRNA positive cells within M1 ipsilateral to the trained paw in

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

SRT and contralateral to the trained paw in the ACT group were also significantly higher than for controls. This most likely reflects the fact that ACT also involves some motor skill learning – although the skill is far less complex than SRT. Similarly, ACT rats show an increase of dendritic branching and spine density in M1 layer V neurons when compared to inactive control animals, but those plastic changes are less pronounced as compared to SRT rats (Wang et al., 2011). Increased levels of *Arc* mRNA in the ipsilateral M1 in SRT as well as the non-significant trend towards a correlation for *Arc* expression and learning success may reflect the contribution of the ipsilateral hemisphere to the learning process (Chen et al., 1997), however an unspecific bystander effect, e.g., through transcallosal activation, cannot be excluded.

That training-induced changes were confined to M1 and were not detectable within RMA, S1 and ST highlights the importance of M1 for motor learning. M1 has been proposed to be the site of storage of motor memory traces (Monfils et al., 2005). Protein synthesis in M1 is required for consolidation of a motor skill (Luft et al., 2004) and motor learning is accompanied by ~~various changes in M1:~~

~~a) At the network level, a transient enlargement of the forelimb representation can be observed after reach training (Kleim et al., 1998, Molina-Luna et al., 2008). The degree of enlargement correlates with performance improvements (Molina-Luna et al., 2008). b) At the cellular level, training stimulates the formation of novel synapses (Kleim et al., 2004) that become selectively preserved over time if they are functionally relevant (Xu et al., 2009). c) On the level of intercellular connectivity, motor training alters synaptic weights.~~

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Feldfunktion geändert

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

~~Synapses of layer II/III horizontal connections become synaptic-strengthening ed~~

via LTP-like mechanisms ~~during motor learning~~ (Rioul-Pedotti et al., 1998, Rioul-Pedotti et al., 2000). It thus seems highly plausible that motor memories are stored within M1 circuitry.

~~The training-induced expression of Arc mRNA in M1 supports this view.~~ Through an interaction with dynamin and endophilin 2 and 3, Arc is the rate-limiting molecule in endocytosis of AMPA-type glutamate receptors at synaptic sites (Bramham et al., 2008). By regulating the homeostatic synaptic scaling of AMPA receptors (Shepherd et al., 2006), Arc is ideally suited to control synaptic strength and cellular excitability (Chowdhury et al., 2006, Rial Verde et al., 2006, Bramham et al., 2008). Moreover, Arc is an essential protein for LTP consolidation (Guzowski et al., 2000, Plath et al., 2006). Arc may not only regulate the translational machinery required for LTP consolidation (Bramham et al., 2010), by controlling F-actin polymerization it also modulates structural changes like the expansion of postsynaptic spines underlying late-phase LTP (Bourne and Harris, 2008). Such an expansion of postsynaptic spine width in combination with LTP in layer 1 horizontal connections has been also observed within M1 of rats that learned a reaching task (Harms et al., 2008).

~~Induction of Arc mRNA that occurs within minutes after a neuron became activated in relation to behavioral experience (Guzowski et al., 1999) seems to be an early step specifically tied to the encoding of a motor memory in M1. However, subsequent cascades of gene expression and maturation processes likely have to take place until the memory is fully consolidated (Luft and Buitrago,~~

Formatiert: Nicht Hervorheben

Formatiert: Nicht Hervorheben

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert

...

~~2005). Arc mRNA trafficking, local translation of Arc mRNA into protein as well as the turnover of Arc mRNA is a strictly regulated process (Giorgi et al., 2007, Huang et al., 2007, Park et al., 2008, Bramham et al., 2010). Thus, the broad expression pattern of Arc mRNA comprising all cortical layers of M1 (except layer I) (Fig. 1b and c) may reflect an initial activation of neuronal populations in response to training. Later different processes may be required to selectively stabilize newly formed synapses (Xu et al., 2009) or strengthen specific synaptic connections that are functionally relevant.~~

In summary, our findings suggest a mechanistic link between M1 Arc expression and motor skill learning in rats. As training-dependent Arc expression was confined to M1 therefore suggests the formation of motor memory traces in M1 neurons.

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Feldfunktion geändert

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Schriftart: Cambria, Schriftartfarbe: Schwarz, Rechtschreibung und Grammatik prüfen

Formatiert: Englisch (USA)

Figure legends

Figure 1. *Arc* mRNA expression within M1. (A) More *Arc* mRNA positive neurons were found in the hemisphere contralateral to the trained forelimb of

SRT animals (cSRT) as compared to the other groups (iSRT: 69%; cACT: 69%; iACT: 66%; control group: 52%; **: $p < 0.001$). *: M1 of control animals (CG) show less *Arc* mRNA expression when compared to the iSRT and the cACT group ($p < 0.05$). Error bars indicate SEM. **(B)** and **(C)**. Representative confocal images taken from primary motor cortex of different groups. *Arc* mRNA positive neurons are distributed across all cortical layers except layer I. Note: sections in B were processed in a different FISH cycle than those shown in C. Vertical bars indicate the borders of cortical layers. Scale bar: 150 μm .

Figure 2. Relationship between *Arc* mRNA expression in M1 and learning success (between sessions 1 and 2). The number of *Arc* mRNA positive neurons in the cSRT group ($n=18$) is significantly correlated to the learning rate ($R=0.523$, $p=0.0259$; grey lines indicate the 95% confidence interval).

Figure 3. *Arc* mRNA expression across different brain regions. Neither for RMA (**A**; $p=0.349$), S1 (**B**; $p=0.7$) nor ST (**C**; $p=0.298$) statistically significant differences were found between groups). Error bars indicate SEM.

Formatiert: Englisch (USA)

Acknowledgements

This work was supported by grants from The Hertie Foundation and the Swiss National Fund (SNF, NCCR Neuro) and a fellowship to B.M.W.-B. of the German

Formatiert: Englisch (USA)

National Academic Foundation. [We thank Ana Fulgencio Maisch and Clement Osei-Atiemo for technical support.](#)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

References

- Bourne JN, Harris KM (2008) Balancing structure and function at hippocampal dendritic spines. Annu Rev Neurosci 31:47-67.
- Bramham CR, Alme MN, Bittins M, Kuipers SD, Nair RR, Pai B, Panja D, Schubert M, Soule J, Tiron A, Wibrand K (2010) The Arc of synaptic memory. Exp Brain Res 200:125-140.
- Bramham CR, Worley PF, Moore MJ, Guzowski JF (2008) The immediate early gene arc/arg3.1: regulation, mechanisms, and function. J Neurosci 28:11760-11767.
- Buitrago MM, Ringer T, Schulz JB, Dichgans J, Luft AR (2004) Characterization of motor skill and instrumental learning time scales in a skilled reaching task in rat. Behav Brain Res 155:249-256.
- Chen R, Gerloff C, Hallett M, Cohen LG (1997) Involvement of the ipsilateral motor cortex in finger movements of different complexities. Ann Neurol 41:247-254.
- Chowdhury S, Shepherd JD, Okuno H, Lyford G, Petralia RS, Plath N, Kuhl D, Huganir RL, Worley PF (2006) Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. Neuron 52:445-459.
- Conner JM, Culbertson A, Packowski C, Chiba AA, Tuszynski MH (2003) Lesions of the Basal forebrain cholinergic system impair task acquisition and abolish cortical plasticity associated with motor skill learning. Neuron 38:819-829.
- Fletcher BR, Calhoun ME, Rapp PR, Shapiro ML (2006) Fornix lesions decouple the induction of hippocampal arc transcription from behavior but not plasticity. J Neurosci 26:1507-1515.
- Giorgi C, Yeo GW, Stone ME, Katz DB, Burge C, Turrigiano G, Moore MJ (2007) The EJC factor eIF4AIII modulates synaptic strength and neuronal protein expression. Cell 130:179-191.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Guzowski JF (2002) Insights into immediate-early gene function in hippocampal memory consolidation using antisense oligonucleotide and fluorescent imaging approaches. Hippocampus 12:86-104.

Formatiert: Englisch (USA)

Guzowski JF, Lyford GL, Stevenson GD, Houston FP, McGaugh JL, Worley PF, Barnes CA (2000) Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory. J Neurosci 20:3993-4001.

Formatiert: Englisch (USA)

Guzowski JF, McNaughton BL, Barnes CA, Worley PF (1999) Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. Nat Neurosci 2:1120-1124.

Formatiert: Englisch (USA)

Hanlon EC, Faraguna U, Vyazovskiy VV, Tononi G, Cirelli C (2009) Effects of skilled training on sleep slow wave activity and cortical gene expression in the rat. Sleep 32:719-729.

Formatiert: Englisch (USA)

Hikosaka O (2002) Central mechanisms of motor skill learning. Curr Opin Neurobiol 12:217-222.

Huang F, Chotiner JK, Steward O (2007) Actin polymerization and ERK phosphorylation are required for Arc/Arg3.1 mRNA targeting to activated synaptic sites on dendrites. J Neurosci 27:9054-9067.

Formatiert: Englisch (USA)

Harms KJ, Rioult-Pedotti MS, Carter DR, Dunaevsky A (2008) Transient spine expansion and learning-induced plasticity in layer 1 primary motor cortex. J Neurosci 28:5686-5690.

Formatiert: Englisch (USA)

Kleim JA, Barbay S, Nudo RJ (1998) Functional reorganization of the rat motor cortex following motor skill learning. J Neurophysiol 80:3321-3325.

Formatiert: Englisch (USA)

Kleim JA, Bruneau R, Calder K, Pocock D, VandenBerg PM, MacDonald E, Monfils MH, Sutherland RJ, Nader K (2003) Functional organization of adult motor cortex is dependent upon continued protein synthesis. Neuron 40:167-176.

Formatiert: Englisch (USA)

Kleim JA, Hogg TM, VandenBerg PM, Cooper NR, Bruneau R, Remple M (2004) Cortical synaptogenesis and motor map reorganization occur during late, but not early, phase of motor skill learning. J Neurosci 24:628-633.

Formatiert: Englisch (USA)

Kleim JA, Lussnig E, Schwarz ER, Comery TA, Greenough WT (1996) Synaptogenesis and Fos expression in the motor cortex of the adult rat after motor skill learning. J Neurosci 16:4529-4535.

Formatiert: Englisch (USA)

Luft AR, Buitrago MM (2005) Stages of motor skill learning. Mol Neurobiol 32:205-216.

Formatiert: Englisch (USA)

Luft AR, Buitrago MM, Ringer T, Dichgans J, Schulz JB (2004) Motor skill learning depends on protein synthesis in motor cortex after training. J Neurosci 24:6515-6520.

Formatiert: Englisch (USA)

Lyford GL, Yamagata K, Kaufmann WE, Barnes CA, Sanders LK, Copeland NG, Gilbert DJ, Jenkins NA, Lanahan AA, Worley PF (1995) Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. Neuron 14:433-445.

Formatiert: Englisch (USA)

McIntyre CK, Miyashita T, Setlow B, Marjon KD, Steward O, Guzowski JF, McGaugh JL (2005) Memory-influencing intra-basolateral amygdala drug infusions modulate expression of Arc protein in the hippocampus. Proc Natl Acad Sci U S A 102:10718-10723.

Formatiert: Englisch (USA)

Miyashita T, Kubik S, Lewandowski G, Guzowski JF (2008) Networks of neurons, networks of genes: an integrated view of memory consolidation. Neurobiol Learn Mem 89:269-284.

Formatiert: Englisch (USA)

Molina-Luna K, Hertler B, Buitrago MM, Luft AR (2008) Motor learning transiently changes cortical somatotopy. Neuroimage 40:1748-1754.

Formatiert: Englisch (USA)

Monfils MH, Plautz EJ, Kleim JA (2005) In search of the motor engram: motor map plasticity as a mechanism for encoding motor experience. Neuroscientist 11:471-483.

Formatiert: Englisch (USA)

Panja D, Dagate G, Bidinosti M, Wibrand K, Kristiansen AM, Sonenberg N, Bramham CR (2009) Novel translational control in Arc-dependent long term potentiation consolidation in vivo. J Biol Chem 284:31498-31511.

Formatiert: Englisch (USA)

Park S, Park JM, Kim S, Kim JA, Shepherd JD, Smith-Hicks CL, Chowdhury S, Kaufmann W, Kuhl D, Ryazanov AG, Haganir RL, Linden DJ, Worley PF (2008) Elongation factor 2 and fragile X mental retardation protein control

the dynamic translation of Arc/Arg3.1 essential for mGluR-LTD. Neuron 59:70-83.

Paxinos G, Watson C (1997) The rat brain in stereotaxic coordinates. Elsevier Academic Press, Amsterdam.

Formatiert: Englisch (USA)

Plath N, Ohana O, Dammermann B, Errington ML, Schmitz D, Gross C, Mao X, Engelsberg A, Mahlke C, Welzl H, Kobalz U, Stawrakakis A, Fernandez E, Waltereit R, Bick-Sander A, Therstappen E, Cooke SF, Blanquet V, Wurst W, Salmen B, Bosl MR, Lipp HP, Grant SG, Bliss TV, Wolfer DP, Kuhl D (2006) Arc/Arg3.1 is essential for the consolidation of synaptic plasticity and memories. Neuron 52:437-444.

Formatiert: Englisch (USA)

Ramirez-Amaya V, Vazdarjanova A, Mikhael D, Rosi S, Worley PF, Barnes CA (2005) Spatial exploration-induced Arc mRNA and protein expression: evidence for selective, network-specific reactivation. J Neurosci 25:1761-1768.

Formatiert: Englisch (USA)

Rial Verde EM, Lee-Osbourne J, Worley PF, Malinow R, Cline HT (2006) Increased expression of the immediate-early gene arc/arg3.1 reduces AMPA receptor-mediated synaptic transmission. Neuron 52:461-474.

Formatiert: Englisch (USA)

Richter JD, Sonenberg N (2005) Regulation of cap-dependent translation by eIF4E inhibitory proteins. Nature 433:477-480.

Formatiert: Englisch (USA)

Riout-Pedotti MS, Friedman D, Donoghue JP (2000) Learning-induced LTP in neocortex. Science 290:533-536.

Formatiert: Englisch (USA)

Riout-Pedotti MS, Friedman D, Hess G, Donoghue JP (1998) Strengthening of horizontal cortical connections following skill learning. Nat Neurosci 1:230-234.

Formatiert: Englisch (USA)

Shepherd JD, Rumbaugh G, Wu J, Chowdhury S, Plath N, Kuhl D, Huganir RL, Worley PF (2006) Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. Neuron 52:475-484.

Formatiert: Englisch (USA)

Smith-Hicks C, Xiao B, Deng R, Ji Y, Zhao X, Shepherd JD, Posern G, Kuhl D, Huganir RL, Ginty DD, Worley PF, Linden DJ (2010) SRF binding to SRE 6.9 in the Arc promoter is essential for LTD in cultured Purkinje cells. Nat Neurosci 13:1082-1089.

Formatiert: Englisch (USA)

Steward O, Wallace CS, Lyford GL, Worley PF (1998) Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. Neuron 21:741-751.

Vazdarjanova A, Ramirez-Amaya V, Insel N, Plummer TK, Rosi S, Chowdhury S, Mikhael D, Worley PF, Guzowski JF, Barnes CA (2006) Spatial exploration induces ARC, a plasticity-related immediate-early gene, only in calcium/calmodulin-dependent protein kinase II-positive principal excitatory and inhibitory neurons of the rat forebrain. J Comp Neurol 498:317-329.

Formatiert: Englisch (USA)

Wang L, Conner JM, Rickert J, Tuszynski MH (2011) Structural plasticity within highly specific neuronal populations identifies a unique parcellation of motor learning in the adult brain. Proc Natl Acad Sci U S A 108:2545-2550.

Formatiert: Englisch (USA)

Wegenast-Braun BM, Fulgencio Maisch A, Eicke D, Radde R, Herzig MC, Staufenbiel M, Jucker M, Calhoun ME (2009) Independent effects of intra- and extracellular Abeta on learning-related gene expression. Am J Pathol 175:271-282.

Formatiert: Englisch (USA)

Whishaw IQ, Pellis SM (1990) The structure of skilled forelimb reaching in the rat: a proximally driven movement with a single distal rotatory component. Behav Brain Res 41:49-59.

Formatiert: Englisch (USA)

Worley PF, Bhat RV, Baraban JM, Erickson CA, McNaughton BL, Barnes CA (1993) Thresholds for synaptic activation of transcription factors in hippocampus: correlation with long-term enhancement. J Neurosci 13:4776-4786.

Formatiert: Englisch (USA)

Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, Tennant K, Jones T, Zuo Y (2009) Rapid formation and selective stabilization of synapses for enduring motor memories. Nature 462:915-919.

Formatiert: Englisch (USA)

Zou Z, Li F, Buck LB (2005) Odor maps in the olfactory cortex. Proc Natl Acad Sci U S A 102:7724-7729.

Formatiert: Englisch (USA)

- Bourne₁ J.N., Harris₁ K.M. (2008). Balancing structure and function at hippocampal dendritic spines. *Annu Rev Neurosci*. 31: 47-67.
- Bramham₁ C.R., Alme₁ M.N., Bittins₁ M., Kuipers₁ S.D., Nair₁ R.R., Pai₁ B., Panja₁ D., Schubert₁ M., Soule₁ J., Tiron₁ A., Wibrand₁ K. (2010). The Arc of synaptic memory. *Exp Brain Res*. 200: 125-140.
- Bramham₁ C.R., Worley₁ P.F., Moore₁ M.J., Guzowski₁ J.F. (2008). The immediate-early gene arc/arg3.1: regulation, mechanisms, and function. *J Neurosci*. 28: 11760-11767.
- Buitrago₁ M.M., Ringer₁ T., Schulz₁ J.B., Dichgans₁ J., Luft₁ A.R. (2004). Characterization of motor skill and instrumental learning time scales in a skilled-reaching task in rat. *Behav Brain Res*. 155: 249-256.
- Chen₁ R., Gerloff₁ C., Hallett₁ M., Cohen₁ L.G. (1997). Involvement of the ipsilateral motor cortex in finger movements of different complexities. *Ann Neurol*. 41: 247-254.
- Chowdhury₁ S., Shepherd₁ J.D., Okuno₁ H., Lyford₁ G., Petralia₁ R.S., Plath₁ N., Kuhl₁ D., Huganir₁ R.L., Worley₁ P.F. (2006). Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. *Neuron*. 52: 445-459.
- Conner₁ J.M., Culbertson₁ A., Packowski₁ C., Chiba₁ A.A., Tuszynski₁ M.H. (2003). Lesions of the Basal forebrain cholinergic system impair task acquisition and abolish cortical plasticity associated with motor skill learning. *Neuron*. 38: 819-829.
- Fletcher₁ B.R., Calhoun₁ M.E., Rapp₁ P.R., Shapiro₁ M.L. (2006). Fornix lesions decouple the induction of hippocampal arc transcription from behavior but not plasticity. *J Neurosci*. 26: 1507-1515.
- Giorgi₁ C., Yeo₁ G.W., Stone₁ M.E., Katz₁ D.B., Burge₁ C., Turrigiano₁ G., Moore₁ M.J. (2007). The EJC factor eIF4AIII modulates synaptic strength and neuronal protein expression. *Cell*. 130: 179-191.
- Guzowski₁ J.F. (2002). Insights into immediate-early gene function in hippocampal memory consolidation using antisense oligonucleotide and fluorescent imaging approaches. *Hippocampus*. 12: 86-104.

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Guzowski, J.F., Lyford, G.L., Stevenson, G.D., Houston, F.P., McGaugh, J.L., Worley, P.F., Barnes, C.A. (2000). Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory. *J Neurosci*, 20, 3993-4001.

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Guzowski, J.F., McNaughton, B.L., Barnes, C.A., Worley, P.F. (1999). Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat Neurosci*, 2, 1120-1124.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Hanlon, E.G., Faraguna, U., Vyazovskiy, V.V., Tononi, G., Cirelli, C. (2009). Effects of skilled training on sleep slow wave activity and cortical gene expression in the rat. *Sleep*, 32, 719-729.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Hikosaka, O. (2002). Central mechanisms of motor skill learning. *Curr Opin Neurobiol*, 12, 217-222.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Huang, F., Chotiner, J.K., Steward, O. (2007). Actin polymerization and ERK phosphorylation are required for Arc/Arg3.1 mRNA targeting to activated synaptic sites on dendrites. *J Neurosci*, 27, 9054-9067.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Harms, K.J., Rioult-Pedotti, M.S., Carter, D.R., Dunaevsky, A. (2008). Transient spine expansion and learning-induced plasticity in layer 1 primary motor cortex. *J Neurosci*, 28, 5686-5690.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Kleim, J.A., Barbay, S., Nudo, R.J. (1998). Functional reorganization of the rat motor cortex following motor skill learning. *J Neurophysiol*, 80, 3321-3325.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Kleim, J.A., Bruneau, R., Calder, K., Pocock, D., VandenBerg, P.M., MacDonald, E., Monfils, M.H., Sutherland, R.J., Nader, K. (2003). Functional organization of adult motor cortex is dependent upon continued protein synthesis. *Neuron*, 40, 167-176.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Kleim, J.A., Hogg, T.M., VandenBerg, P.M., Cooper, N.R., Bruneau, R., Rempel, M. (2004). Cortical synaptogenesis and motor map reorganization occur during late, but not early, phase of motor skill learning. *J Neurosci*, 24, 628-633.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

- Kleim, J.A., Lussnig, E., Schwarz, E.R., Comery, T.A., Greenough, W.T. (1996). Synaptogenesis and Fos expression in the motor cortex of the adult rat after motor skill learning. *J Neurosci*, 16, 4529-4535.
- Luft, A.R., Buitrago, M.M. (2005). Stages of motor skill learning. *Mol Neurobiol*, 32, 205-216.
- Luft, A.R., Buitrago, M.M., Ringer, T., Dichgans, J., Schulz, J.B. (2004). Motor skill learning depends on protein synthesis in motor cortex after training. *J Neurosci*, 24, 6515-6520.
- Lyford, G.L., Yamagata, K., Kaufmann, W.E., Barnes, C.A., Sanders, L.K., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Lanahan, A.A., Worley, P.F. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron*, 14, 433-445.
- McIntyre, C.K., Miyashita, T., Setlow, B., Marjon, K.D., Steward, O., Guzowski, J.F., McGaugh, J.L. (2005). Memory-influencing intra-basolateral amygdala drug infusions modulate expression of Arc protein in the hippocampus. *Proc Natl Acad Sci U S A*, 102, 10718-10723.
- Miyashita, T., Kubik, S., Lewandowski, G., Guzowski, J.F. (2008). Networks of neurons, networks of genes: an integrated view of memory consolidation. *Neurobiol Learn Mem*, 89, 269-284.
- Molina-Luna, K., Hertler, B., Buitrago, M.M., Luft, A.R. (2008). Motor learning transiently changes cortical somatotopy. *Neuroimage*, 40, 1748-1754.
- Monfils, M.H., Plautz, E.J., Kleim, J.A. (2005). In search of the motor engram: motor map plasticity as a mechanism for encoding motor experience. *Neuroscientist*, 11, 471-483.
- Panja, D., Dwyer, G., Bidinosti, M., Wibrand, K., Kristiansen, A.M., Sonenberg, N., Bramham, C.R. (2009). Novel translational control in Arc-dependent long-term potentiation consolidation in vivo. *J Biol Chem*, 284, 31498-31511.
- Park, S., Park, J.M., Kim, S., Kim, J.A., Shepherd, J.D., Smith-Hicks, C.L., Chowdhury, S., Kaufmann, W., Kuhl, D., Ryazanov, A.G., Huganir, R.L.,

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

- Linden, D.J., Worley, P.F. (2008). Elongation factor 2 and fragile X mental retardation protein control the dynamic translation of Arc/Arg3.1 essential for mGluR-LTD. *Neuron*, 59, 70-83.
- Plath, N., Ohana, O., Dammermann, B., Errington, M.L., Schmitz, D., Gross, C., Mao, X., Engelsberg, A., Mahlke, C., Welzl, H., Kobalz, U., Stawrakakis, A., Fernandez, E., Waltereit, R., Bick-Sander, A., Therstappen, E., Cooke, S.F., Blanquet, V., Wurst, W., Salmen, B., Bosl, M.R., Lipp, H.P., Grant, S.G., Bliss, T.V., Wolfert, D.P., Kuhl, D. (2006). Arc/Arg3.1 is essential for the consolidation of synaptic plasticity and memories. *Neuron*, 52, 437-444.
- Ramirez-Amaya, V., Vazdarjanova, A., Mikhael, D., Rosi, S., Worley, P.F., Barnes, C.A. (2005). Spatial exploration induced Arc mRNA and protein expression: evidence for selective, network-specific reactivation. *J Neurosci*, 25, 1761-1768.
- Rial Verde, E.M., Lee-Osbourne, J., Worley, P.F., Malinow, R., Cline, H.T. (2006). Increased expression of the immediate-early gene arc/arg3.1 reduces AMPA receptor-mediated synaptic transmission. *Neuron*, 52, 461-474.
- Richter, J.D., Sonenberg, N. (2005). Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature*, 433, 477-480.
- Rioult-Pedotti, M.S., Friedman, D., Donoghue, J.P. (2000). Learning-induced LTP in neocortex. *Science*, 290, 533-536.
- Rioult-Pedotti, M.S., Friedman, D., Hess, G., Donoghue, J.P. (1998). Strengthening of horizontal cortical connections following skill learning. *Nat Neurosci*, 1, 230-234.
- Shepherd, J.D., Rumbaugh, G., Wu, J., Chowdhury, S., Plath, N., Kuhl, D., Huganir, R.L., Worley, P.F. (2006). Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. *Neuron*, 52, 475-484.
- Smith-Hicks, C., Xiao, B., Deng, R., Ji, Y., Zhao, X., Shepherd, J.D., Posern, G., Kuhl, D., Huganir, R.L., Ginty, D.D., Worley, P.F., Linden, D.J. (2010).
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)
- Formatiert: Schriftart: Helvetica, Englisch (USA), Rechtschreibung und Grammatik nicht prüfen
- Formatiert: Schriftart: Kursiv
- Formatiert: Schriftart: Kursiv
- Formatiert: Englisch (USA)

- SRF binding to SRE 6.9 in the Arc promoter is essential for LTD in cultured Purkinje cells. *Nat Neurosci*, 13, 1082-1089.
- Steward, O., Wallace, C.S., Lyford, G.L., Worley, P.F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron*, 21, 741-751.
- Vazdarjanova, A., Ramirez-Amaya, V., Insel, N., Plummer, T.K., Resi, S., Chowdhury, S., Mikhael, D., Worley, P.F., Guzowski, J.F., Barnes, C.A. (2006). Spatial exploration induces ARC, a plasticity-related immediate-early gene, only in calcium/calmodulin-dependent protein kinase II-positive principal excitatory and inhibitory neurons of the rat forebrain. *J Comp Neurol*, 498, 317-329.
- Wang, L., Conner, J.M., Rickert, J., Tuszynski, M.H. (2011). Structural plasticity within highly specific neuronal populations identifies a unique parcellation of motor learning in the adult brain. *Proc Natl Acad Sci U S A*, 108, 2545-2550.
- Wegenast-Braun, B.M., Fulgencio-Maisch, A., Eicke, D., Radde, R., Herzig, M.C., Staufenbiel, M., Jucker, M., Calhoun, M.E. (2009). Independent effects of intra- and extracellular Abeta on learning-related gene expression. *Am J Pathol*, 175, 271-282.
- Whishaw, I.Q., Pollis, S.M. (1990). The structure of skilled forelimb reaching in the rat: a proximally driven movement with a single distal rotatory component. *Behav Brain Res*, 41, 49-59.
- Worley, P.F., Bhat, R.V., Baraban, J.M., Erickson, C.A., McNaughton, B.L., Barnes, C.A. (1993). Thresholds for synaptic activation of transcription factors in hippocampus: correlation with long-term enhancement. *J Neurosci*, 13, 4776-4786.
- Xu, T., Yu, X., Perlik, A.J., Tobin, W.F., Zweig, J.A., Tennant, K., Jones, T., Zuo, Y. (2009). Rapid formation and selective stabilization of synapses for enduring motor memories. *Nature*, 462, 915-919.
- Zou, Z., Li, F., Buck, L.B. (2005). Odor maps in the olfactory cortex. *Proc Natl Acad Sci U S A*, 102, 7724-7729.

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Englisch (USA)

